



Plant biology and pathology/Biologie et pathologie végétales

Leaving the meristem behind: The genetic and molecular control of leaf patterning and morphogenesis

Au-delà du méristème : le contrôle génétique et moléculaire de la morphogénèse foliaire

Alice Hasson¹, Thomas Blein^{1,2}, Patrick Laufs^{1,*}

Institut Jean-Pierre-Bourgin, Inra, route de Saint Cyr, 78026 Versailles cedex, France

ARTICLE INFO

Article history:

Available online 12 March 2010

Keywords:

Morphogenesis
Genetic network
Polarity
Signalling

Mots clés :

Morphogénèse
Réseau génétique
Polarité
Signalisation

ABSTRACT

Leaves, which play an essential role in plant photosynthesis, share common features such as being flat structures, but also show an impressive variability in their sizes and shapes. Following its initiation in the meristems, leaf development is patterned along three polarization axes to establish its basic architecture. This process is further complicated in the case of compound leaves with the formation of new growth axes. Growth and differentiation must be properly coordinated to regulate the size and the flatness of the leaf. This review provides an overview of the genetic and molecular regulatory networks underlying leaf development, with an emphasis on leaf polarity and the comparison of simple and compound leaves.

© 2010 Published by Elsevier Masson SAS on behalf of Académie des sciences.

R É S U M É

Les feuilles sont des organes essentiels pour la photosynthèse des plantes et possèdent des caractéristiques communes comme le fait d'être des structures planes, mais montrent aussi une impressionnante variabilité de leurs formes et tailles. Après son initiation au niveau du méristème, la feuille se développe suivant trois axes de polarisation, lui permettant ainsi d'atteindre son architecture de base. La formation de nouveaux axes de croissance lorsqu'il s'agit de feuilles composées rend ce processus plus complexe. La croissance et la différenciation doivent être correctement coordonnées afin de réguler la taille et la planéité de la feuille. Cette revue apporte une vue d'ensemble des réseaux génétiques et moléculaires de régulation du développement foliaire, et met l'accent sur la polarité foliaire et la comparaison entre feuilles simples et composées.

© 2010 Publié par Elsevier Masson SAS pour l'Académie des sciences.

1. Introduction

Leaves are determinate organs that are the main photosynthetic structures of land plants. As such, they

must optimise their light-capturing surface and their gas exchanges with the environment, which would ideally require a leaf as large and thin as possible. However, leaves are also subjected to the physical laws that constrain increases in size and decreases in thickness. Thin leaves are more subject to water loss and heat up more rapidly when exposed to light. Large leaves require more support tissues, which results in a higher biomass investment per unit leaf area [1]. Broad leaves also heat rapidly, especially when the ambient airflow is low [2]. Increasing the level of leaf dissection, via the formation of lobes, helps to cool down

* Corresponding author.

E-mail address: laufs@versailles.inra.fr (P. Laufs).

¹ <http://www.ijpb.versailles.inra.fr/en/bc/equipes/Meristeme2/index.html>

² Present address: Albert-Ludwigs-Universität Freiburg, Institut für Biologie II Botanik, Sonnenstrasse 5, 79104 Freiburg, Germany.

the leaf [2], but also has opposite effects on the surface area available for light capture. On the one hand, it reduces the surface area of each individual leaf, whereas on the other hand, it may increase light interception at the whole plant level by reducing the self-shading of lower leaves by upper leaves [3]. Therefore, the effects of leaf shape and size on plant fitness is a complex issue. For instance, comparison of near-isogenic cotton lines with contrasted leaf shapes showed that leaves with intermediate lobing were the most efficient at the plant level [4]. Such complex, and sometimes opposite constraints may explain the incredible diversity in shapes and sizes that is observed for leaves in different species or even within a single individual, depending on environmental factors or developmental stage (leaf heteroblasty).

In the wide diversity of leaf shapes, two main groups can be distinguished according to the degree of complexity: simple leaves and compound leaves (Fig. 1), [5]. Here, we will present genetic, molecular and hormonal networks acting during leaf initiation, outgrowth and shaping, and present some ideas about how these networks could have been modified to generate contrasting leaf shapes. This review will concentrate on the dicot leaf, either on simple leaves (like the plant model *Arabidopsis thaliana*) or on compound leaves (such as tomato and pea).

2. Leaf initiation at the meristem, or the first step towards determinacy

2.1. The SAM is the ultimate source of all aerial organs

The shoot apical meristem (SAM) contains a pool of stem cells and initiates all the leaves or flowers. The earliest event associated with organ initiation is the switch from an indeterminate to a determinate fate in the founder cells, a small group of cells that will give rise to the lateral

organs. The mechanisms controlling meristem function or the process of primordium initiation will not be detailed here, as recent reviews are available, including in this issue [6–12], but we will highlight some of the specific points that are essential to understand later steps of leaf development.

2.2. Leaving the meristem behind: the switch from indeterminate to determinate fate during primordium initiation

The SAM is characterized by the expression of Class I *KNOTTED1-LIKE HOMEODOMAIN (KNOXI)* genes represented in *Arabidopsis* by the *SHOOT MERISTEMLESS (STM)*, *BREVIPEDICELLUS/KNOTTED-LIKE FROM Arabidopsis1 (BP/KNAT1)*, *KNAT2* and *KNAT6* genes [13–17]. These homeodomain transcription factors are required for the establishment and maintenance of the meristems, in part via the activation of the cytokinin pathway [13,18]. They show distinct expression patterns in the SAM but share a common zone of repression at the site of the incipient leaf primordium (Fig. 2) [13,19]. This repression marks the transition to a determinate fate and is essential for proper leaf development, as ectopic *KNOXI* expression during simple leaf development leads to severe defects, including leaf lobing and formation of ectopic meristems [20–22].

Given the importance of proper regulation of *KNOXI* genes, it is not surprising that several pathways contribute to their repression in the leaf. One of these pathways involves auxin. Local auxin accumulation, resulting from PIN1-mediated polar transport of this hormone, determines the site of primordium initiation [6,8,12,23]. Real-time imaging of developing apices and genetic analysis showed that these peaks of auxin also contribute to *KNOXI* repression (Fig. 2) [24,25]. Another pathway repressing *KNOXI* genes involves the *ARP* genes coding for MYB-domain




Fig. 1. Overview of the different leaf architectures. Two main leaf architectures are recognized: simple leaves formed by a single blade supported by the petiole (A–C), and compound leaves with several units called leaflets (D, E). Compound leaves are further divided into pinnate leaves when the leaflets are attached on different parts of an elongated axis called the rachis (D), and into palmate leaves when all leaflets unit on a single point (E). The leaf or leaflet margin is either entire (A), serrated (B, D, E) or lobed (C). Leaves are organised along three axes: proximodistal (petiole to blade tip), adaxial-abaxial (or dorsoventral from the upper to the lower side) and mediolateral (midrib to margin). Shadows of common European trees leaves are shown: magnolia (A), hornbeam (B, F), maple (C), ash (D) and chestnut tree (E).




Fig. 2. Genetic and molecular network controlling leaf initiation and differentiation. *KNOXI* genes are expressed in the SAM, where they repress the *ARP* genes and the GA pathway to maintain cells in an undifferentiated state. Local auxin accumulation sets up the site of primordium initiation, and together with the *ARP* genes, represses *KNOXI* expression. Repression of the *KNOXI* genes in the primordium allows the GA pathway to be derepressed, thus leading to growth and differentiation. *KNOXI* contribute to the expression of the *GA 2-oxidase* that inactivates GA.

transcription factors, and named after *Arabidopsis* *ASYMETRIC LEAVES1* (*AS1*), maize *ROUGH SHEATH2* (*RS2*) and *Antirrhinum majus* *PHANTASTICA* (*PHAN*) genes [26–29]. *ARP* genes are specifically expressed in lateral organ founder cells where they repress *KNOXI* expression. In turn, *AS1* expression is excluded from the meristem by *STM*, leading to a complementary expression pattern of *KNOXI* and *ARP* genes (Fig. 2) [26]. The mechanism of *AS1*-mediated *KNOXI* repression was recently elucidated [30]. *AS1* forms a repressor complex, together with the LOB domain protein *AS2*, the predicted RNA binding protein *RIK* and the chromatin-remodeling protein *HIRA*. This complex binds to the promoter of the *BP* and *KNAT2* genes during early stages of primordium development and represses the expression of these two *KNOXI* genes. Such a repression was proposed to occur via chromatin modifications that lead to a silenced state that is stably transmitted through cell divisions during later stages of leaf development [30].

Besides morphological modifications, ectopic *KNOXI* expression in leaves also leads to a cellular phenotype characterized by reduced cell expansion and differentiation [10]. Such a phenotype suggests a defect in the gibberellin (GA) pathway, a hypothesis supported by the observation that normal leaf development is restored in *KNOXI* overexpressors by the activation of the GA pathway [31]. Indeed, recent evidence suggests that a large part of

KNOXI protein function is mediated by its inhibitory effects on GA synthesis and signalling [8,32]. First, *KNOXI* transcription factors of different species directly down-regulate the expression of the *GA 20-OXIDASE* gene that codes for an enzyme controlling a key step of GA biosynthesis (Fig. 2) [31,33,34]. Furthermore, *STM* promotes the expression of the *GA 2-OXIDASE* gene involved in the inactivation of GA, which protects the meristem from GAs that may diffuse from the primordium [35]. Thus, the combined effects of *KNOXI* inhibition of GA synthesis and promotion of GA inactivation lead to reduced GA signalling. Such *KNOXI*-mediated repression of GA signalling is required for maintaining the meristem in an undifferentiated state, but when this pathway is ectopically activated in the leaf, it delays cell expansion and differentiation and leads to abnormal leaf development.

2.3. Defining the organ boundaries

Proper lateral organ development also requires the physical separation of individual organ primordia, which is accomplished through the action of the three *Arabidopsis* *CUP-SHAPED COTYLEDON1*, 2 and 3 (*CUC1*, 2 and 3) genes. These genes encode NO APICAL MERISTEM/ATAF/CUP-SHAPED COTYLEDON (*NAC*) plant-specific transcription factors and are specifically expressed in a narrow file of cells at the base of organs that marks their boundary with neighbouring organs or with the meristem (Fig. 2) [36–38]. This specific expression pattern may be a response to the local peaks of auxin associated with primordium formation [25]. Inactivation of one or several of the *CUC* genes leads to various levels of organ fusion [36–38]. *CUC1* and *CUC2*, are post-transcriptionally regulated by a microRNA, miR164, like all other members of the *NAM* clade, named after the petunia NO APICAL MERISTEM (*NAM*) gene [39]. In contrast, *CUC3* is not targeted by miR164. miRNAs are part of a diverse family of small regulatory non-coding RNAs and have emerging roles in the regulation of plant development [40,41]. Indeed, miR164-mediated regulation of *CUC1/CUC2* is important for the homeostasis of the boundary domain [42–46].

3. Building the basic leaf shape: how and why are the polarization axes setup?

3.1. A functional polarisation of the leaf

Emerging from the SAM, the leaf primordium is a radial structure that will rapidly organize itself along three axes: proximodistal, mediolateral and adaxial-abaxial (or dorsoventral) (Figs. 1 and 3A). Here, we will only discuss the mechanism of leaf polarization along the adaxial-abaxial axis, as it is the best understood from a molecular and genetic perspective, and is the most relevant for the physiology of the leaf. Early polarization along the adaxial-abaxial axis during primordium development serves to specify cell types within the mature leaf. The adaxial side of the leaf blade is exposed to the sun and differentiates into palisade cells that are optimised for space filling and have a high chloroplast content, thus maximizing light interception. The opposite abaxial side is constituted of




Fig. 3. Adaxial-abaxial leaf polarity and leaf architecture. A. Adaxial-abaxial leaf polarity and meristem. (1) Scheme showing the orientation of the adaxial and abaxial domains of a young primordium relative to the shoot apical meristem; and (2) a transverse section of an expanded leaf. B. Adaxial-abaxial leaf polarity, blade expansion and meristem formation. (3) A mature leaf shows an expanded blade formed by adaxial and abaxial domains and supports a secondary meristem in its axillary adaxial region; (4) Adaxialized mutants such as *phb-1D* form leaves with no blade expansion that carry ectopic meristems on their lower base; (5) Abaxialized mutants such as *kan* mutants develop bladeless leaves that lack meristems; (6) The *Antirrhinum phan* mutant with patches of abaxial tissues on its adaxial side and ectopic outgrowths; (7) Ectopic meristems form on the adaxial side of the blade of plants overexpressing *KNOXI* genes. C. Adaxial-abaxial leaf polarity and signalling from the meristem. (8) An incision made between the meristem and a young primordium leads to the formation of a radial abaxialized organ. (9) If the incision is made later, once the primordium is already polarized, a leaf with a normal organization develops (10). These experiments provide evidence for a meristem-derived signal that promotes adaxial fate and/or represses abaxial fate.

spongy parenchymatous cells leaving intercellular spaces that promote gas exchanges. Stomata distribution is also often polarized, with a higher density on the abaxial side. Finally, vascular tissues are also polarized along this axis, as the xylem is only found in the adaxial side and the phloem in the abaxial side.

The genetic control of adaxial-abaxial leaf polarity involves a multilayered regulatory network entailing a complex interplay between several classes of transcription factors and other regulators subjected to different levels of regulation. Furthermore, establishing this polarity is not an autonomous process occurring only in the leaf primordium but also involves communication with the meristem.

3.2. The antagonism between adaxial and abaxial factors controls leaf polarization

The core regulatory module controlling adaxial-abaxial leaf polarity is based on the antagonistic relationship between two sets of transcription factors that determine the identity of the adaxial and abaxial leaf domains. The class III *HOMEODOMAIN-LEUCINE ZIPPER* (*HD-ZIPIII*) genes *REVOLUTA* (*REV*), *PHABULOSA* (*PHB*) and *PHAVOLUTA* (*PHV*) of *Arabidopsis* are expressed on the adaxial side of the primordium [47], whereas members of the *KANADI* (*KAN*) and *YABBY* (*YAB*) gene families are expressed on the abaxial side (Fig. 4) [48,49]. These genes determine the identity of the leaf domain in which they are expressed, and repress the expression of the identity genes of the complementary domain (Fig. 4). For instance, leaves of *kan* loss-of-function

mutants show an adaxialized phenotype (Fig. 3B) and an expansion of the expression of the *HD-ZIPIII* genes, whereas on the contrary, *KAN* over-expression leads to leaf abaxialization and *HD-ZIPIII* repression [50]. In addition to this antagonism between adaxial and abaxial factors, *YAB* expression is strongly reduced by the inactivation of the *KAN* genes, suggesting that the *YAB* genes act downstream of the *KAN* (Fig. 4) [50].

3.3. Several partially redundant pathways contribute to reinforce polarization along the adaxial-abaxial axis

Interacting with this core-regulatory module, several other factors control leaf polarity. Most of these factors appear to modify the activity of the *HD-ZIPIII*, *KAN* and *YAB* components. Examples of such factors include the *LITTLE ZIPPER* (*ZPR*) proteins [51,52]. The *ZPR* proteins form heterodimers with *HD-ZIPIII* and prevent their binding to DNA. Interestingly, *ZPR* expression is induced by *HD-ZIPIII*, thus establishing a negative feedback loop (Fig. 4). This modulation of the activity of the *HD-ZIPIII* proteins appears, however, to have only a small contribution to leaf polarity, in contrast to the pathways described below.

Whereas the *phan* mutant revealed a central contribution of this gene to *Antirrhinum* adaxial-abaxial leaf polarity (Fig. 3B), the initially identified *Arabidopsis as1* and *as2* mutants showed no obvious polarity defect. However, identification of novel *as1/as2* alleles in a different genetic background [53], and further genetic studies revealed the contribution of the *AS1/AS2* pathway




Fig. 4. Genetic and molecular network controlling adaxial-abaxial leaf polarity. Two main genes families are involved in the definition of the identity of the adaxial and abaxial leaf domains: the *KAN* and *HD-ZIPIII* genes are, respectively, expressed in the abaxial and the adaxial domains, and thus define the identity of each territory. The antagonism between these two groups is at the root of their complementary expression domains in the leaf. The contribution of several other molecular actors reinforces these expression patterns: *miR165/166* negatively regulates *HD-ZIPIII* genes, while *ARP* proteins promote *HD-ZIPIII* expression. In parallel, *ARP* proteins negatively control *KAN* and *miR165/166* expressions. *KAN* factors activate the expression of the *YAB* genes that contribute to define the abaxial identity, while *YAB* genes are subjected to negative regulation by the *ARP* and *TAS3* pathways.

for the promotion of adaxial leaf fate [54–57]. *AS1*, together with *AS2*, positively regulate *HD-ZIPIII* expression, while they negatively regulate *KAN* and *YAB* expression (Fig. 4) [54–57]. If the *AS1/AS2* pathway regulates components of the core regulatory module, the inverse also occurs: *KAN1* directly represses *AS2* expression in the abaxial domain of the leaf (Fig. 4) [58]. This reveals another mutually repressive loop between the adaxial *AS1/AS2* factors and the abaxial *KAN* determinants, which contributes to the proper separation of the adaxial and abaxial domains. Finally, mutations in several ribosomal proteins or in components of the proteasome enhance the leaf polarity defects of *as1* or *as2* mutants [59–61]. However, the mechanism by which the regulation of protein translation and turnover feeds into the control of leaf polarity remains unknown.

Members of two classes of small RNAs are also involved in leaf polarity control. The related *miR165/miR166* miRNAs have a prominent role in abaxial leaf fate. These miRNAs target the *HD-ZIPIII* *PHB*, *REV* and *PHV* transcripts (Fig. 4). All gain-of-function mutants of these genes identified thus far are mutated in the *miR165/166* binding site [62,63], suggesting that miRNA-dependent regulation of these genes is critical for normal plant development. However, as these mutations also disrupt a StAR-related lipid transfer (*START*) domain, known to interact with lipids in animal systems, it was not clear whether the phenotype associated with these mutations was a consequence of a defective *miR165/miR166* regulation, a modification of the properties of the *START* domain or a combination of both. The last two hypotheses were

excluded by the observation that the expression of *miR165/166*-insensitive versions of *HD-ZIPIII* genes that possess silent mutations in the miRNA binding site (and therefore code for unmodified proteins) could phenocopy the original gain-of-function alleles [62,64]. The *miR165/166* was reported to be expressed in the abaxial domain [65] or throughout the leaf primordium [56] and contributes to restrict *HD-ZIPIII* expression to the adaxial domain [65]. Interestingly, some evidence suggests that the *AS1/AS2* pathway may increase *HD-ZIPIII* activity by reducing *miR165/166* expression, consistent with the repressing activity of the *AS1/AS2* complex [56].

The second class of small RNAs provides a link between leaf polarity and auxin. Auxin response is mediated by the AUXIN RESPONSE FACTORS (*ARF*), a class of transcription factors that binds to auxin-responsive promoter elements (*ARE*) in target genes regulated by auxin [66]. Two of these *ARFs* genes, *ARF3/ETTIN* and *ARF4*, were identified following a screen for suppressors of *KAN* overexpressors, thus providing the first genetic evidence for the involvement of auxin in adaxial-abaxial polarity. The expression of these two genes overlaps in the abaxial domain (Fig. 4), and their simultaneous inactivation leads to organ adaxialization [67]. Further genetic studies revealed, however, that the *ARF3/ARF4* are not just merely downstream of the *KAN* pathway, but are more likely to cooperate with *KAN* for the promotion of abaxial fate. *ARF3* and *ARF4* are regulated by trans-acting small interfering RNAs (*ta-siRNAs*) produced from the *TAS3* gene [54,68–70]. Although mutating components involved in *ta-siRNA* biogenesis or expressing a *TAS3*-insensitive *ARF3* gene has an effect on leaf heteroblasty, no obvious effects on leaf polarity were observed [69,70]. However, such defects are observed when mutations of the *ta-siRNA* biogenesis machinery are combined with *as1* mutations [40]. At the molecular level, the *AS1* and *TAS3* pathways redundantly repress the expression of *YAB* genes. This indicates a functional redundancy between the *TAS3* and *AS1* factors in the promotion of adaxial identity and the repression of abaxial fate.

3.4. Setting up polarity by mobile signals

Whereas the complex regulatory network described above accounts for the maintenance of the two complementary leaf domains, it cannot explain the establishment of polarity nor the alignment of leaf polarity with respect to the axis of the plant. Early expression studies suggested that the leaf initium is not polarized, as mRNA of both adaxial and abaxial determinants were detected throughout the initium [47,71]. However, recent advancements in real-time imaging of developing lateral organs indicate that polarization is concomitant with the formation of the initium [25]. Production of a diffusible signal is a mechanism widely used in animal systems to polarize organs or tissues, and recent evidence suggests that movement of *ta-siRNA* could play a similar role during leaf polarization. Due to a localized expression of the *TAS3* gene and/or components of the *ta-siRNA* production machinery, *TAS3 ta-siRNA* are produced only in the outermost cells layers of the adaxial domain of the

primordium [72–74]. The TAS3 ta-siRNAs diffuse across a limited number of cells, generating a gradient over the adaxial-abaxial axis. Computer simulations indicate that this gradient may sharpen the expression pattern of the *ARF3/4* genes [75]. Interestingly, in maize the distribution of one component of the ta-siRNA production machinery remains polar in a radialized leaf of a mutant in which the expression of other genes becomes apolar [76]. This suggests that, at least in the maize leaf, ta-siRNA gradients have a prominent role in ad/abxial polarization.

Leaf polarization is always coordinated with respect to the meristem. Indeed, forward-looking microsurgical experiments made more than 50 years ago showed the existence of a signal emanating from the meristem and polarising the leaf. Incisions separating incipient primordia from the SAM result in radially abaxialized structures, when later incisions did not modify an already polarized organ (Fig. 3C) [77], suggesting that the signal is needed only to establish the polarity and not for its maintenance. Recent re-examination of these experiments further showed that an intact epidermal layer is required for meristem-primodium signalling [78]. What is the signal? Currently, this question is still open, but a candidate has arisen. It was proposed that a sterol/lipid molecule produced by the meristem could form a gradient and interact with the HD-ZIPIII and modify its activity [47]. Since the adaxial and abaxial domains would be exposed to different concentrations of the signal, a polarity could be initiated. No further observations however support this hypothesis, and the signal has not yet been identified.

3.5. Abaxial/adaxial polarization is important for all aspects of leaf function

Besides determining proper cell types, adaxial-abaxial polarization is also important for two other aspects of leaf development. First, proper definition of the two adaxial and abaxial domains is important for leaf flattening and blade outgrowth. Adaxialized or abaxialized mutants form finger-like leaves with no blade expansion (Fig. 3B) [50,79]. Furthermore, in some leaves of the *Antirrhinum phan* mutant, patches of abaxial tissues appear on the adaxial side of the leaf, and ridges resembling blades initiate from the boundary between these regions with different identities (Fig. 3B) [29]. These observations led to the hypothesis that signalling across the juxtaposed abaxial and adaxial domains is necessary to promote blade outgrowth [29]. The molecular mechanisms that mediate this signalling and promote blade outgrowth are currently unknown. However, genetic analysis points to a role of the *YAB* genes in the promotion of blade outgrowth [50]. Alternatively, a recent study underscored the role of elongated, specialized cells located at the margin of the leaf blade (the margin cells) and the possible role of brassinolides in the control of leaf growth and shaping [80]. Whether this is relevant for early blade outgrowth or limited to later stages of leaf shaping remains an open question.

The second implication of the adaxial-abaxial polarization is the formation of axillary meristems. These structures are normally located in the axil, i.e. on the adaxial side of the junction of the leaf petiole with the

stem, and ectopic meristems that form on the leaf blade of *KNOX1* overexpressors, for instance, are also always located on the adaxial side (Fig. 3B) [22]. Furthermore, meristem formation is perturbed in leaf polarity mutants. Adaxialized leaves develop axillary meristems from the lower part of the leaf base [79], whereas abaxialized mutants are defective for meristem formation (Fig. 3B) [62]. Altogether, this demonstrates a link between the adaxial identity and the ability to develop meristems. The nature of this link is not yet understood, but some determinants of the adaxial domain such as *PHB* are also expressed in the meristem, suggesting that they could provide a molecular link between meristem function and adaxial leaf fate [47,81].

3.6. The same players but different roles? Conservation and specificities of the adaxial-abaxial regulatory network in different species

Except for some species (such as most cacti) in which the leaves are transformed into spines [82], flattening and polarization along the adaxial-abaxial axis is a general feature of most leaves. Although the knowledge of the regulatory networks at work in this process is by far best understood in *Arabidopsis*, more and more data become available for other species, such as the dicot *Antirrhinum* or the monocots maize and rice. The general picture that emerges from this overview is that most of the proteins or RNA regulators described above show a high level of conservation within vascular plants [83]; for instance, the maize *RS2 ARP* gene fully complements the *Arabidopsis as1* mutant [84]. Although the function may be conserved, the importance of each pathway can be significantly different between species. An example of this is the different importance of the *ARP* pathway in *Arabidopsis* and *Antirrhinum* leaf polarity, as mentioned above. Another example is the different contributions of the *TAS3* pathway, as witnessed by the difference in the severity of the leaf polarity defects observed in *Arabidopsis* and maize following the inactivation of the *SGS3* gene that codes for a component of the ta-siRNA production machinery [54,68,73]. Therefore, the regulatory network may show significant differences between species, perhaps resulting from different developmental or environmental constraints.

4. Shaping the mature leaf: patterning and growth control

Once the three axes of polarization are set up, further patterning will occur in the leaf primordium to shape the vascular network and the leaf margin. Positional information provided by epidermal foci of auxin accumulation and routes of auxin flow from the margin to the centre of the primordium pattern the vasculature in the developing leaf [85]. In parallel, the epidermal auxin maxima determine the position of the marginal outgrowths that form the serrations [24]. A factor possibly downstream of auxin that mediates leaf margin dissection is the *CUC2* gene. This gene is expressed in the sinus of the developing serrations and its activity, controlled by the miRNA miR164, determines

the level of leaf dissection [86]. This dissecting role is conserved for *CUC2* orthologues in several Eudicots [87,88]. All together, this indicates that the auxin/*CUC*/miR164 regulatory module acting during primordium initiation is also used during leaf margin sculpting.

In the subsequent phases of development, the leaf primordium gains in size and reaches its mature shape. Initially, cells present a high proliferation activity throughout the blade. Then, a proliferation arrest front progresses in a basipetal way, from the tip of the leaf to its base. After proliferation arrest, growth is achieved via cell expansion [89]. Whereas the basic components of the cell machinery have been identified and largely characterized, how their function is integrated during leaf development to generate a flat structure with a given shape and size remains unclear. Many mutants developing abnormal leaf shapes have been characterized over the years, and the defects of certain of these mutants have been traced back to variations in cell proliferation or expansion patterns [90–92]. A useful example of this is the *Antirrhinum CINCINNATA (CIN)* gene, which encodes a TEOSINTE BRANCHED/CYCLOIDEA/PROLIFERATING CELL FACTOR (TCP) transcription factor and which controls cell proliferation arrest. In *cin* mutants, leaves are crinkly, as a result of excess growth at their margins [93]. In *Arabidopsis*, a similar phenotype is found in the *jaw-D* dominant mutant, where the *MIRJAW (MIR319)* gene that targets several *CIN*-related *TCPs* is over-expressed [94].

5. Making compound leaves in a complex way

5.1. Delaying differentiation to elaborate compound leaves

Whereas *KNOXI* genes are repressed during the entire process of simple leaf development, these genes are activated again during the development of most

compound leaves following a transient phase of repression (Figs. 5 and 6) [95,96]. Recent work on an emerging model for compound leaf development, the *Arabidopsis* relative *Cardamine hirsuta*, showed that *KNOXI* activity is necessary for compound leaf formation (Fig. 5B) [97]. Reciprocally, a higher or prolonged *KNOXI* activity increases the complexity of compound leaves. For instance, ectopic expression of *KNOXI* transgenes leads to higher-order leaflets in cardamine and tomato (Fig. 5B) [96,97]. However, the activity of the *KNOXI* proteins is also modified indirectly by alteration in their protein interaction network. *KNOXI* proteins rely for their nuclear localization and DNA binding on the formation of heterodimers with another class of homeoproteins, the *BEL* proteins [11]. Furthermore, the *BEL* proteins interact with a newly identified truncated *KNOXI* protein called *PTS/TDK1* [98]. Increasing the expression of the *PTS/TDK1* gene or inactivating a specific *BEL* member leads to tomato leaves of higher complexity [98]. Interestingly, an increased expression of the *PTS/TDK1* gene, due to a single nucleotide deletion in its promoter, is the basis for the increased leaf complexity of a wild tomato species collected by Charles Darwin from the Galapagos Islands [98]. All together, these observations indicate that *KNOXI* activity acts like a rheostat to control compound leaf complexity. How is this achieved? Based on its known role in the meristem, reactivation of *KNOXI* gene expression during compound leaf development may delay cell differentiation, and, thus, create a pseudomeristematic environment that is required for the elaboration of complex leaf structures. As in the meristem, *KNOXI* maintains an undifferentiated state by repressing the *GA* pathway in compound leaf primordia. Exogenous application of *GA*, or increased *GA* signalling due to the inactivation of the *DELLA*-type gene *PROCERA*, simplifies tomato leaf architecture and partly antagonizes the effects of *KNOXI* overexpression [31,99]. All together, this reveals




Fig. 5. Compound leaf development. A. Compound leaf development. (1–4) The initial developmental steps of a compound leaf are similar to those of a simple leaf: *KNOXI* genes are repressed in the initiation and then, are turned-on again to maintain an undifferentiated state. Leaflet primordia are initiated in a process resembling the initiation of a leaf primordium. B. Compound leaf architecture modification. (5) Scheme of a normal compound leaf bearing several primary leaflets. (6) *KNOXI* genes overexpression leads to the formation of ectopic, higher order leaflets. (7) On the contrary, inactivation of *KNOXI* genes transforms the compound leaf into a simple structure. (8) Perturbation of auxin transport/response or reduced activity of the *NAM/CUC3* boundary genes leads to fewer leaflets that can be partially fused.




Fig. 6. Genetic and molecular network controlling leaflet formation. Leaflet formation requires the maintenance of an undifferentiated environment to which contribute the KNOXI, LFY-like and TCP pathways. Action of the KNOXI factors is mediated by an inhibition of the Gibberellin (GA) pathway, while the TCP genes are negatively regulated by the microRNA, miR319. The site of leaflet initiation is determined by local accumulation of auxin (sun diagram). *NAM/CUC3* genes are expressed in the distal boundary of the incipient leaflet (represented in red), where they locally repress cell proliferation and thus lead to the formation of groove. *NAM* genes are negatively regulated by the microRNA, miR164. *NAM/CUC3* genes contribute to leaflet formation, possibly via a positive feed-forward loop with the KNOXI and LFY-like pathways.

the recruitment of the meristematic KNOXI/GA regulatory module to the developing primordium of compound leaves (Figs. 5 and 6).

In addition, to the KNOXI/GA pathway, tomato leaflet formation is also controlled by the *TCP* gene *LANCEOLATE* (*LA*). The expression of this gene gradually builds-up during leaf development, and may, like its *Arabidopsis* orthologue *CIN*, trigger growth arrest and differentiation. Precocious expression in the semi-dominant *Lanceolate* (*La*) allele that carries a mutation in the miR319 binding site leads to the formation of simple tomato leaves [100]. It is suggested that the *LA* gene defines a developmental window during which leaflet initiation is possible.

Despite the widespread association of *KNOXI* expression with compound leaf development [95], an alternate pathway has been revealed for some species. The initial observation that *KNOXI* expression is absent from developing compound primordia of pea [101] was recently enlarged to several members of a *Fabaceae* sub-clade that, in addition to the pea, includes fava pea and alfalfa, and that diverged 39 million years ago from other *Fabaceae* [102]. What is controlling leaflet initiation and formation in species of this clade? The answer to this question came from the identification of the gene affected in the pea *unifoliata* (*uni*) mutant that develops simplified leaves. *UNI* codes for the pea orthologue of the *LEAFY* (*LFY*) and *FLORICAULA* (*FLO*) genes that are required to specify the identity of the floral meristem [103,104]. In the pea, *UNI* has conserved its role in the control of floral development

but, in addition, it is also required for maintaining leaf primordia cells in an undifferentiated state to allow compound leaf formation (Fig. 6). The contribution of *LFY* orthologues during compound leaf development has been shown for alfalfa that, like the pea, does not express *KNOXI* genes in their leaves [105], and also in the tomato, a species that expresses *KNOXI* genes during leaf development [106]. This suggests that the *KNOXI* and *LFY* pathways cooperate to enable compound leaf development, and that the relative contribution of each of these pathways varies depending on the species [107].

5.2. Organizing the outgrowth of new structures

Whereas the two pathways described above may explain the higher organogenetic potential of compound leaves, other factors are required to organize growth into the new structures that will form the leaflets. First, auxin, whose role in leaf primordium initiation and marginal serrations has been described above, is also involved during compound leaf formation (Figs. 5 and 6). More precisely, peaks of auxin response contribute to the outgrowth of leaflet primordia, whereas repression of the auxin response is required to inhibit outgrowth of the regions between leaflets (Fig. 5B) [108,109]. Second, orthologues of the *Arabidopsis* *CUC* genes are required for proper individualization of the leaflets. These genes show a conserved expression pattern in the axils of the outgrowing leaflets and their inactivation leads to leaflet fusions (Fig. 5B) [87,88]. In addition, inactivation of these genes leads to a reduction in the number of leaflets, thus revealing a promoting effect of the *CUC* genes on leaflet formation, a role that could be based on a positive feedback loop between the *CUC* genes and the *KNOXI* and/or *LFY* pathways (Fig. 6) [88].

5.3. Simple leaves versus compound leaves

The relationship between simple and compound leaves has long been a matter of debate [5]. Two opposite hypotheses have been proposed. The first one suggests that each leaflet of a compound leaf is homologous to a simple leaf, and that a compound leaf is equivalent to a compressed shoot. The second hypothesis suggests that leaflets result from extreme dissections of the leaf margins, and that the entire compound leaf is homologous to a simple leaf. Does the recent knowledge gained in leaf development help to discriminate between these hypotheses? During compound leaf development, the redeployment of the *CUC*/auxin/*KNOXI*/GA regulatory modules that are active in the meristem supports the hypothesis of a shoot-like identity of compound leaves. This identity, however, is only partial as compound leaf primordia do not express stem cell markers characteristic of the meristem [97]. Alternatively, serrations can be transformed into leaflets [97,110], and the auxin/*CUC* module is involved in both serration and leaflet formation, supporting the homology between these two structures. Taken together, these observations suggest that instead of a unique model for all compound leaves, these structures may arise through different developmental paths that converge to

generate mature structures with comparable architectures. The repeated independent apparition of compound leaves during evolution also supports this idea [95].

6. Conclusion

Leaf development, from its initiation to growth and differentiation, is an excellent model to address fundamental issues of biology such as the generation of complex forms and their evolution. During the last years, important progress has been made in the identification and functional analysis of genetic and molecular regulators of leaf development. This research has revealed a surprisingly limited set of players, that can either play the same game with slightly different rules (during adaxial-abaxial leaf polarization, for instance), or play different games with similar rules (during the formation of different compound leaves, for instance). Understanding the differences in the rules, and how they modify the behaviour of the players to generate different games is the next challenge.

Acknowledgements

The authors thank Anne Plessis, Elizabeth F. Crowell and Michael J. Harrington for their helpful comments on the manuscript.

References

- [1] U. Niinemets, A. Portsmuth, M. Tobias, Leaf shape and venation pattern alter the support investments within leaf lamina in temperate species: a neglected source of leaf physiological differentiation? *Funct. Ecol.* 21 (2007) 28–40.
- [2] S. Vogel, Leaves in the lowest and highest winds: temperature, force and shape, *New Phytol.* 183 (2009) 13–26.
- [3] M.S. Ali, K. Kikuzawa, Shoot morphology of *Aucuba japonica* incurred by anisophylly: ecological implications, *J. Plant. Res.* 118 (2005) 329–338.
- [4] R. Wells, W.R. Meredith, J.R. Williford, Canopy photosynthesis and its relationship to plant productivity in near-isogenic cotton lines differing in leaf morphology, *Plant Physiol.* 82 (1986) 635–640.
- [5] C. Champagne, N. Sinha, Compound leaves: equal to the sum of their parts? *Development* 131 (2004) 4401–4412.
- [6] A.J. Fleming, The control of leaf development, *New Phytol.* 166 (2005) 9–20.
- [7] M.R. Tucker, T. Laux, Connecting the paths in plant stem cell regulation, *Trends Cell. Biol.* 17 (2007) 403–410.
- [8] B. Veit, Hormone mediated regulation of the shoot apical meristem, *Plant Mol. Biol.* 69 (2009) 397–408.
- [9] N. Carraro, A. Peaucelle, P. Laufs, J. Traas, Cell differentiation and organ initiation at the shoot apical meristem, *Plant Mol. Biol.* 60 (2006) 811–826.
- [10] S. Scofield, J.A. Murray, KNOX gene function in plant stem cell niches, *Plant Mol. Biol.* 60 (2006) 929–946.
- [11] O. Hamant, V. Pautot, Plant Development: a TALE story, *C. R. Biologies* 333 (2010) 371–381.
- [12] I. Bohn-Courseau, Auxin: a major regulator of organogenesis, *C. R. Biologies* 333 (2010) 290–296.
- [13] J.A. Long, E.I. Moan, J.I. Medford, M.K. Barton, A member of the KNOTTED class of homeodomain proteins encoded by the STM gene of *Arabidopsis*, *Nature* 379 (1996) 66–69.
- [14] S.J. Douglas, G. Chuck, R.E. Dengler, L. Peleacanda, C.D. Riggs, KNAT1 and ERECTA regulate inflorescence architecture in *Arabidopsis*, *Plant Cell* 14 (2002) 547–558.
- [15] S.P. Venglat, T. Dumonceaux, K. Rozwadowski, L. Parnell, V. Babic, W. Keller, R. Martienssen, G. Selvaraj, R. Datla, The homeobox gene BREVIPEDICELLUS is a key regulator of inflorescence architecture in *Arabidopsis*, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 4730–4735.
- [16] E. Belles-Boix, O. Hamant, S.M. Witiak, H. Morin, J. Traas, V. Pautot, KNAT6: an *Arabidopsis* homeobox gene involved in meristem activity and organ separation, *Plant Cell* 18 (2006) 1900–1907.
- [17] V. Pautot, J. Dockx, O. Hamant, J. Kronenberger, O. Grandjean, D. Jublot, J. Traas, KNAT2: evidence for a link between knotted-like genes and carpel development, *Plant Cell* 13 (2001) 1719–1734.
- [18] E. Vollbrecht, L. Reiser, S. Hake, Shoot meristem size is dependent on inbred background and presence of the maize homeobox gene, knotted1, *Development* 127 (2000) 3161–3172.
- [19] D. Jackson, B. Veit, S. Hake, Expression of maize *KNOTTED1* related homeobox genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot, *Development* 120 (1994) 405–413.
- [20] C. Lincoln, J. Long, J. Yamaguchi, K. Serikawa, S. Hake, A knotted1-like homeobox gene in *Arabidopsis* is expressed in the vegetative meristem and dramatically alters leaf morphology when overexpressed in transgenic plants, *Plant Cell* 6 (1994) 1859–1876.
- [21] N. Ori, Y. Eshed, G. Chuck, J.L. Bowman, S. Hake, Mechanisms that control knox gene expression in the *Arabidopsis* shoot, *Development* 127 (2000) 5523–5532.
- [22] N.R. Sinha, R.E. Williams, S. Hake, Overexpression of the maize homeobox gene, KNOTTED-1, causes a switch from determinate to indeterminate cell fates, *Genes Dev.* 7 (1993) 787–795.
- [23] R. Benjamins, B. Scheres, Auxin: the looping star in plant development, *Annu. Rev. Plant Biol.* 59 (2008) 443–465.
- [24] A. Hay, M. Barkoulas, M. Tsiantis, ASYMMETRIC LEAVES1 and auxin activities converge to repress BREVIPEDICELLUS expression and promote leaf development in *Arabidopsis*, *Development* 133 (2006) 3955–3961.
- [25] M.G. Heisler, C. Ohno, P. Das, P. Sieber, G.V. Reddy, J.A. Long, E.M. Meyerowitz, Patterns of auxin transport and gene expression during primordium development revealed by live imaging of the *Arabidopsis* inflorescence meristem, *Curr. Biol.* 15 (2005) 1899–1911.
- [26] M.E. Byrne, R. Barley, M. Curtis, J.M. Arroyo, M. Dunham, A. Hudson, R.A. Martienssen, Asymmetric leaves1 mediates leaf patterning and stem cell function in *Arabidopsis*, *Nature* 408 (2000) 967–971.
- [27] M.C. Timmermans, A. Hudson, P.W. Bercraft, T. Nelson, ROUGH SHEATH2: a Myb protein that represses knox homeobox genes in maize lateral organ primordia, *Science* 284 (1999) 151–153.
- [28] M. Tsiantis, R. Schneeberger, J.F. Golz, M. Freeling, J.A. Langdale, The maize rough sheath2 gene and leaf development programs in monocot and dicot plants, *Science* 284 (1999) 154–156.
- [29] R. Waites, A. Hudson, Phantastica: a gene required for dorsoventrality of leaves in *Antirrhinum majus*, *Development* 121 (1995) 2143–2154.
- [30] M. Guo, J. Thomas, G. Collins, M.C. Timmermans, Direct repression of KNOX loci by the ASYMMETRIC LEAVES1 complex of *Arabidopsis*, *Plant Cell* 20 (2008) 48–58.
- [31] A. Hay, H. Kaur, A. Phillips, P. Hedden, S. Hake, M. Tsiantis, The gibberellin pathway mediates KNOTTED1-type homeobox function in plants with different body plans, *Curr. Biol.* 12 (2002) 1557–1565.
- [32] E. Shani, O. Yanai, N. Ori, The role of hormones in shoot apical meristem function, *Curr. Opin. Plant Biol.* 9 (2006) 484–489.
- [33] H. Chen, A.K. Banerjee, D.J. Hannapel, The tandem complex of BEL and KNOX partners is required for transcriptional repression of *ga20ox1*, *Plant J.* 38 (2004) 276–284.
- [34] T. Sakamoto, N. Kamiya, M. Ueguchi-Tanaka, S. Iwahori, M. Matsuoka, KNOX homeodomain protein directly suppresses the expression of a gibberellin biosynthetic gene in the tobacco shoot apical meristem, *Genes Dev.* 15 (2001) 581–590.
- [35] S. Jasinski, P. Piazza, J. Craft, A. Hay, L. Woolley, I. Rieu, A. Phillips, P. Hedden, M. Tsiantis, KNOX action in *Arabidopsis* is mediated by coordinate regulation of cytokinin and gibberellin activities, *Curr. Biol.* 15 (2005) 1560–1565.
- [36] M. Aida, T. Ishida, H. Fukaki, H. Fujisawa, M. Tasaka, Genes involved in organ separation in *Arabidopsis*: an analysis of the cup-shaped cotyledon mutant, *Plant Cell* 9 (1997) 841–857.
- [37] S. Takada, K. Hibara, T. Ishida, M. Tasaka, The CUP-SHAPED COTYLEDON1 gene of *Arabidopsis* regulates shoot apical meristem formation, *Development* 128 (2001) 1127–1135.
- [38] C.W. Vroemen, A.P. Mordhorst, C. Albrecht, M.A. Kwaaitaal, S.C. de Vries, The CUP-SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in *Arabidopsis*, *Plant Cell* 15 (2003) 1563–1577.
- [39] E. Souer, A. Van Houwelingen, D. Kloos, J. Mol, R. Koes, The *No Apical Meristem* gene of petunia is required for pattern formation in embryos and flower and is expressed at meristem and primordia boundaries, *Cell* 85 (1996) 159–170.
- [40] D. Garcia, A miRacle in plant development: role of microRNAs in cell differentiation and patterning, *Semin. Cell. Dev. Biol.* 19 (2008) 586–595.

- [41] O. Voinnet, Origin, biogenesis, and activity of plant microRNAs, *Cell* 136 (2009) 669–687.
- [42] C.C. Baker, P. Sieber, F. Wellmer, E.M. Meyerowitz, The early extra petals1 mutant uncovers a role for microRNA miR164c in regulating petal number in *Arabidopsis*, *Curr. Biol.* 15 (2005) 303–315.
- [43] P. Laufs, A. Peaucelle, H. Morin, J. Traas, MicroRNA regulation of the CUC genes is required for boundary size control in *Arabidopsis* meristems, *Development* 131 (2004) 4311–4322.
- [44] A.C. Mallory, D.V. Dugas, D.P. Bartel, B. Bartel, MicroRNA regulation of NAC-domain targets is required for proper formation and separation of adjacent embryonic, vegetative, and floral organs, *Curr. Biol.* 14 (2004) 1035–1046.
- [45] A. Peaucelle, H. Morin, J. Traas, P. Laufs, Plants expressing a miR164-resistant CUC2 gene reveal the importance of post-meristematic maintenance of phyllotaxy in *Arabidopsis*, *Development* 134 (2007) 1045–1050.
- [46] P. Sieber, F. Wellmer, J. Gheyselinck, J.L. Riechmann, E.M. Meyerowitz, Redundancy and specialization among plant microRNAs: role of the MIR164 family in developmental robustness, *Development* 134 (2007) 1051–1060.
- [47] J.R. McConnell, J. Emery, Y. Eshed, N. Bao, J. Bowman, M.K. Barton, Role of PHABULOSA and PHAVOLUTA in determining radial patterning in shoots, *Nature* 411 (2001) 709–713.
- [48] Y. Eshed, S.F. Baum, J.V. Perea, J.L. Bowman, Establishment of polarity in lateral organs of plants, *Curr. Biol.* 11 (2001) 1251–1260.
- [49] R.A. Kerstetter, K. Bollman, R.A. Taylor, K. Bomblied, R.S. Poethig, KANADI regulates organ polarity in *Arabidopsis*, *Nature* 411 (2001) 706–709.
- [50] Y. Eshed, A. Izhaki, S.F. Baum, S.K. Floyd, J.L. Bowman, Asymmetric leaf development and blade expansion in *Arabidopsis* are mediated by KANADI and YABBY activities, *Development* 131 (2004) 2997–3006.
- [51] Y.S. Kim, S.G. Kim, M. Lee, I. Lee, H.Y. Park, P.J. Seo, J.H. Jung, E.J. Kwon, S.W. Suh, K.H. Paek, C.M. Park, HD-ZIP III activity is modulated by competitive inhibitors via a feedback loop in *Arabidopsis* shoot apical meristem development, *Plant Cell* 20 (2008) 920–933.
- [52] S. Wenkel, J. Emery, B.H. Hou, M.M. Evans, M.K. Barton, A feedback regulatory module formed by LITTLE ZIPPER and HD-ZIP III genes, *Plant Cell* 19 (2007) 3379–3390.
- [53] L. Xu, Y. Xu, A. Dong, Y. Sun, L. Pi, H. Huang, Novel as1 and as2 defects in leaf adaxial-abaxial polarity reveal the requirement for ASYMMETRIC LEAVES1 and 2 and ERECTA functions in specifying leaf adaxial identity, *Development* 130 (2003) 4097–4107.
- [54] D. Garcia, S.A. Collier, M.E. Byrne, R.A. Martienssen, Specification of leaf polarity in *Arabidopsis* via the trans-acting siRNA pathway, *Curr. Biol.* 16 (2006) 933–938.
- [55] H. Iwakawa, M. Iwasaki, S. Kojima, Y. Ueno, T. Soma, H. Tanaka, E. Semiarti, Y. Machida, C. Machida, Expression of the ASYMMETRIC LEAVES2 gene in the adaxial domain of *Arabidopsis* leaves represses cell proliferation in this domain and is critical for the development of properly expanded leaves, *Plant J.* 51 (2007) 173–184.
- [56] H. Li, L. Xu, H. Wang, Z. Yuan, X. Cao, Z. Yang, D. Zhang, Y. Xu, H. Huang, The Putative RNA-dependent RNA polymerase RDR6 acts synergistically with ASYMMETRIC LEAVES1 and 2 to repress BREVIDICELLUS and MicroRNA165/166 in *Arabidopsis* leaf development, *Plant Cell* 17 (2005) 2157–2171.
- [57] W.C. Lin, B. Shuai, P.S. Springer, The *Arabidopsis* LATERAL ORGAN BOUNDARIES-domain gene ASYMMETRIC LEAVES2 functions in the repression of KNOX gene expression and in adaxial-abaxial patterning, *Plant Cell* 15 (2003) 2241–2252.
- [58] G. Wu, W.C. Lin, T. Huang, R.S. Poethig, P.S. Springer, R.A. Kerstetter, KANADI1 regulates adaxial-abaxial polarity in *Arabidopsis* by directly repressing the transcription of ASYMMETRIC LEAVES2, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 16392–16397.
- [59] V. Piron, J.P. Etchells, P. Rossignol, S.A. Collier, J.M. Arroyo, R.A. Martienssen, M.E. Byrne, Three PIGGYBACK genes that specifically influence leaf patterning encode ribosomal proteins, *Development* 135 (2008) 1315–1324.
- [60] Y. Yao, Q. Ling, H. Wang, H. Huang, Ribosomal proteins promote leaf adaxial identity, *Development* 135 (2008) 1325–1334.
- [61] W. Huang, L. Pi, W. Liang, B. Xu, H. Wang, R. Cai, H. Huang, The proteolytic function of the *Arabidopsis* 26S proteasome is required for specifying leaf adaxial identity, *Plant Cell* 18 (2006) 2479–2492.
- [62] J.F. Emery, S.K. Floyd, J. Alvarez, Y. Eshed, N.P. Hawker, A. Izhaki, S.F. Baum, J.L. Bowman, Radial patterning of *Arabidopsis* shoots by class III HD-ZIP and KANADI genes, *Curr. Biol.* 13 (2003) 1768–1774.
- [63] M.W. Rhoades, B.J. Reinhart, L.P. Lim, C.B. Burge, B. Bartel, D.P. Bartel, Prediction of plant microRNA targets, *Cell* 110 (2002) 513–520.
- [64] A.C. Mallory, B.J. Reinhart, M.W. Jones-Rhoades, G. Tang, P.D. Zamore, M.K. Barton, D.P. Bartel, MicroRNA control of PHABULOSA in leaf development: importance of pairing to the microRNA 5' region, *EMBO J.* 23 (2004) 3356–3364.
- [65] C.A. Kidner, R.A. Martienssen, Spatially restricted microRNA directs leaf polarity through ARGONAUTE1, *Nature* 428 (2004) 81–84.
- [66] W.D. Teale, I.A. Paponov, K. Palme, Auxin in action: signalling, transport and the control of plant growth and development, *Nat. Rev. Mol. Cell. Biol.* 7 (2006) 847–859.
- [67] I. Pekker, J.P. Alvarez, Y. Eshed, Auxin response factors mediate *Arabidopsis* organ asymmetry via modulation of KANADI activity, *Plant Cell* 17 (2005) 2899–2910.
- [68] X. Adenot, T. Elmayan, D. Laressergues, S. Boutet, N. Bouche, V. Gascioli, H. Vaucheret, DRB4-dependent TAS3 trans-acting siRNAs control leaf morphology through AGO7, *Curr. Biol.* 16 (2006) 927–932.
- [69] N. Fahlgren, T.A. Montgomery, M.D. Howell, E. Allen, S.K. Dvorak, A.L. Alexander, J.C. Carrington, Regulation of AUXIN RESPONSE FACTOR3 by TAS3 ta-siRNA affects developmental timing and patterning in *Arabidopsis*, *Curr. Biol.* 16 (2006) 939–944.
- [70] C. Hunter, M.R. Willmann, G. Wu, M. Yoshikawa, M. de la Luz Gutierrez-Nava, S.R. Poethig, Trans-acting siRNA-mediated repression of ETTIN and ARF4 regulates heteroblasty in *Arabidopsis*, *Development* 133 (2006) 2973–2981.
- [71] K.R. Siegfried, Y. Eshed, S.F. Baum, D. Otsuga, G.N. Drews, J.L. Bowman, Members of the YABBY gene family specify abaxial cell fate in *Arabidopsis*, *Development* 126 (1999) 4117–4128.
- [72] D.H. Chitwood, F.T. Nogueira, M.D. Howell, T.A. Montgomery, J.C. Carrington, M.C. Timmermans, Pattern formation via small RNA mobility, *Genes Dev.* 23 (2009) 549–554.
- [73] F.T. Nogueira, S. Madi, D.H. Chitwood, M.T. Juarez, M.C. Timmermans, Two small regulatory RNAs establish opposing fates of a developmental axis, *Genes Dev.* 21 (2007) 750–755.
- [74] R. Schwab, A. Maizel, V. Ruiz-Ferrer, D. Garcia, M. Bayer, M. Crespi, O. Voinnet, R.A. Martienssen, Endogenous TasIRNAs mediate non-cell autonomous effects on gene regulation in *Arabidopsis thaliana*, *PLoS One* 4 (2009) e5980.
- [75] E. Levine, P. McHale, H. Levine, Small regulatory RNAs may sharpen spatial expression patterns, *PLoS Comput. Biol.* 3 (2007) e233.
- [76] F.T. Nogueira, D.H. Chitwood, S. Madi, K. Ohtsu, P.S. Schnable, M.J. Scanlon, M.C. Timmermans, Regulation of small RNA accumulation in the maize shoot apex, *PLoS Genet.* 5 (2009) e1000320.
- [77] I.M. Sussex, Experiments on the cause of dorsoventrality in leaves, *Nature* 174 (1954) 351–352.
- [78] D. Reinhardt, M. Frenz, T. Mandel, C. Kuhlemeier, Microsurgical and laser ablation analysis of leaf positioning and dorsoventral patterning in tomato, *Development* 132 (2005) 15–26.
- [79] J.R. McConnell, M.K. Barton, Leaf polarity and meristem formation in *Arabidopsis*, *Development* 125 (1998) 2935–2942.
- [80] B. Reinhardt, E. Hanggi, S. Muller, M. Bauch, J. Wyrzykowska, R. Kerstetter, S. Poethig, A.J. Fleming, Restoration of DWF4 expression to the leaf margin of a *dwf4* mutant is sufficient to restore leaf shape but not size: the role of the margin in leaf development, *Plant J.* 52 (2007) 1094–1104.
- [81] S.P. Grigg, C. Canales, A. Hay, M. Tsiantis, SERRATE coordinates shoot meristem function and leaf axial patterning in *Arabidopsis*, *Nature* 437 (2005) 1022–1026.
- [82] J.D. Mauseth, Structure-function relationships in highly modified shoots of cactaceae, *Ann. Bot. (Lond)* 98 (2006) 901–926.
- [83] S.K. Floyd, J.L. Bowman, The ancestral developmental tool kit of land plants, *Int. J. Plant. Sci.* 168 (2007) 1–35.
- [84] G. Theodoris, N. Inada, M. Freeling, Conservation and molecular dissection of ROUGH SHEATH2 and ASYMMETRIC LEAVES1 function in leaf development, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 6837–6842.
- [85] E. Scarpella, D. Marcos, J. Friml, T. Berleth, Control of leaf vascular patterning by polar auxin transport, *Genes Dev.* 20 (2006) 1015–1027.
- [86] K. Nikovics, T. Blein, A. Peaucelle, T. Ishida, H. Morin, M. Aida, P. Laufs, The balance between the MIR164A and CUC2 genes controls leaf margin serration in *Arabidopsis*, *Plant Cell* 18 (2006) 2929–2945.
- [87] Y. Berger, S. Harpaz-Saad, A. Brand, H. Melnik, N. Sirding, J.P. Alvarez, M. Zinder, A. Samach, Y. Eshed, N. Ori, The NAC-domain transcription factor GOBLET specifies leaflet boundaries in compound tomato leaves, *Development* 136 (2009) 823–832.
- [88] T. Blein, A. Pulido, A. Viallette-Guiraud, K. Nikovics, H. Morin, A. Hay, I.E. Johansen, M. Tsiantis, P. Laufs, A conserved molecular framework for compound leaf development, *Science* 322 (2008) 1835–1839.
- [89] P.M. Donnelly, D. Bonetta, H. Tsukaya, R.E. Dengler, N.G. Dengler, Cell cycling and cell enlargement in developing leaves of *Arabidopsis*, *Dev. Biol.* 215 (1999) 407–419.
- [90] E. Anastasiou, M. Lenhard, Growing up to one's standard, *Curr. Opin. Plant Biol.* 10 (2007) 63–69.
- [91] L. De Veylder, T. Beeckman, D. Inze, The ins and outs of the plant cell cycle, *Nat. Rev. Mol. Cell. Biol.* 8 (2007) 655–665.

- [92] H. Tsukaya, Mechanism of leaf shape determination, *Annu. Rev. Plant Biol.* 57 (2006) 477–496.
- [93] U. Nath, B.C. Crawford, R. Carpenter, E. Coen, Genetic control of surface curvature, *Science* 299 (2003) 1404–1407.
- [94] J.F. Palatnik, E. Allen, X. Wu, C. Schommer, R. Schwab, J.C. Carrington, D. Weigel, Control of leaf morphogenesis by microRNAs, *Nature* 425 (2003) 257–263.
- [95] G. Bharathan, T.E. Goliber, C. Moore, S. Kessler, T. Pham, N.R. Sinha, Homologies in leaf form inferred from KNOX1 gene expression during development, *Science* 296 (2002) 1858–1860.
- [96] D. Hareven, T. Gutfinger, A. Parnis, Y. Eshed, E. Lifschitz, The making of a compound leaf: genetic manipulation of leaf architecture in tomato, *Cell* 84 (1996) 735–744.
- [97] A. Hay, M. Tsiantis, The genetic basis for differences in leaf form between *Arabidopsis thaliana* and its wild relative *Cardamine hirsuta*, *Nat. Genet.* 38 (2006) 942–947.
- [98] S. Kimura, D. Koenig, J. Kang, F.Y. Yoong, N. Sinha, Natural variation in leaf morphology results from mutation of a novel KNOX gene, *Curr. Biol.* 18 (2008) 672–677.
- [99] S. Jasinski, A. Tattersall, P. Piazza, A. Hay, J.F. Martinez-Garcia, G. Schmitz, K. Theres, S. McCormick, M. Tsiantis, PROCERA encodes a DELLA protein that mediates control of dissected leaf form in tomato, *Plant J.* 56 (2008) 603–612.
- [100] N. Ori, A.R. Cohen, A. Etzioni, A. Brand, O. Yanai, S. Shleizer, N. Menda, Z. Amsellem, I. Efroni, I. Pekker, J.P. Alvarez, E. Blum, D. Zamir, Y. Eshed, Regulation of LANCEOLATE by miR319 is required for compound-leaf development in tomato, *Nat. Genet.* 39 (2007) 787–791.
- [101] J. Hofer, C. Gourelay, A. Michael, T.H. Ellis, Expression of a class 1 knotted1-like homeobox gene is down-regulated in pea compound leaf primordia, *Plant Mol. Biol.* 45 (2001) 387–398.
- [102] C.E. Champagne, T.E. Goliber, M.F. Wojciechowski, R.W. Mei, B.T. Townsley, K. Wang, M.M. Paz, R. Geeta, N.R. Sinha, Compound leaf development and evolution in the legumes, *Plant Cell* 19 (2007) 3369–3378.
- [103] J. Hofer, L. Turner, R. Hellens, M. Ambrose, P. Matthews, A. Michael, N. Ellis, UNIFOLIATA regulates leaf and flower morphogenesis in pea, *Curr. Biol.* 7 (1997) 581–587.
- [104] E. Moyroud, G. Tichtinsky, F. Parcy, The LEAFY floral regulators in Angiosperms: conserved proteins with diverse roles, *J. Plant Biol.* 52 (2009) 177–185.
- [105] H. Wang, J. Chen, J. Wen, M. Tadege, G. Li, Y. Liu, K.S. Mysore, P. Ratet, R. Chen, Control of compound leaf development by FLORICAULA/LEAFY ortholog SINGLE LEAFLET1 in *Medicago truncatula*, *Plant Physiol.* 146 (2008) 1759–1772.
- [106] N. Molinero-Rosales, M. Jamilena, S. Zurita, P. Gomez, J. Capel, R. Lozano, FALSIFLORA, the tomato orthologue of FLORICAULA and LEAFY, controls flowering time and floral meristem identity, *Plant J.* 20 (1999) 685–693.
- [107] T. Blein, A. Hasson, P. Laufs, Leaf development: what it needs to be complex, *Curr. Opin. Plant Biol.* 13 (2009) 75–82.
- [108] M. Barkoulas, A. Hay, E. Kougioumoutzi, M. Tsiantis, A developmental framework for dissected leaf formation in the *Arabidopsis* relative *Cardamine hirsuta*, *Nat. Genet.* 40 (2008) 1136–1141.
- [109] D. Koenig, E. Bayer, J. Kang, C. Kuhlemeier, N. Sinha, Auxin patterns *Solanum lycopersicum* leaf morphogenesis, *Development* 136 (2009) 2997–3006.
- [110] S. Barth, T. Geier, K. Eimert, B. Watillon, R.S. Sangwan, S. Gleissberg, KNOX overexpression in transgenic *Kohleria* (Gesneriaceae) prolongs the activity of proximal leaf blastozones and drastically alters segment fate, *Planta* 230 (2009) 1081–1091.